



National
Aeronautics and
Space
Administration

NASA Scientific and Technical Document Availability Authorization (DAA)

The DAA approval process applies to all forms of published NASA Scientific and Technical Information (STI), whether disseminated in print or electronically. It is to be initiated by the responsible NASA Project Officer, Technical Monitor, author, or other appropriate NASA official for all presentations, reports, papers, and proceedings that contain NASA STI. Explanations are on the back of this form and are presented in greater detail in NPG 2200.2, "Guidelines for Documentation, Approval, and Dissemination of NASA Scientific and Technical Information."

☐ Original
☐ Modified

I. DOCUMENT/PROJECT IDENTIFICATION

TITLE Wading Bird Populations of the Kennedy Space Center.		AUTHOR(S) Smith, R. B. and D.R. Breininger.
ORIGINATING NASA ORGANIZATION Kennedy Space Center Biomedical Office		PERFORMING ORGANIZATION (If different) Dynamac Corporation, KSC
CONTRACT/GRANT/INTERAGENCY/PROJECT NUMBER(S)	DOCUMENT NUMBER(S) Bullentin of Marine Sci. 57: 230-236	DOCUMENT DATE 1995

For presentations, documents, or other STI to be externally published (including through electronic media), enter appropriate information on the intended publication such as name, place, and date of conference, periodical, or journal name, or book title and publisher in the next box. These documents must be routed to the NASA Headquarters or Center Export Control Administrator for approval (see Sections III and VIII).



II. SECURITY CLASSIFICATION

CHECK ONE (One of the five boxes denoting Security Classification must be checked.)

☐ SECRET ☐ SECRET RD ☐ CONFIDENTIAL ☐ CONFIDENTIAL RD ☒ UNCLASSIFIED

III. AVAILABILITY CATEGORY

<input type="checkbox"/> ITAR <input type="checkbox"/> EAR	Export Controlled Document - USML Category Classification Number (ECCN) <i>(Documents marked in this block must have the concurrence/approval of the NASA Headquarters or Center Export Control Administrator (see Section VIII).)</i>
<input type="checkbox"/> TRADE SECRET <input type="checkbox"/> SBIR <input type="checkbox"/> COPYRIGHTED	Confidential Commercial Document (check appropriate box at left and indicate below the appropriate limitation and expiration): <input type="checkbox"/> U.S. Government agencies and U.S. Government agency contractors only <input type="checkbox"/> NASA contractors and U.S. Government only <input type="checkbox"/> U.S. Government agencies only <input type="checkbox"/> NASA personnel and NASA contractors only <input type="checkbox"/> NASA personnel only <input type="checkbox"/> Available only with the approval of issuing office: <input type="checkbox"/> Limited until (date)
<input checked="" type="checkbox"/> PUBLICLY AVAILABLE	Publicly available documents must be unclassified, may not be export controlled, may not contain trade secret or confidential commercial data, and should have cleared any applicable patents application process.

IV. DOCUMENT DISCLOSING AN INVENTION

THIS DOCUMENT MAY BE RELEASED ON (date)	NASA HQ OR CENTER PATENT OR INTELLECTUAL PROPERTY COUNSEL SIGNATURE	DATE
---------------------------------------------------	----------------------------------------------------------------------------	-------------

V. BLANKET RELEASE (OPTIONAL)

<input type="checkbox"/> All documents issued under the following contract/grant/project number may be processed as checked in Sections II and III.
<input type="checkbox"/> The blanket release authorization granted on (date) <input type="checkbox"/> is RESCINDED - Future documents must have individual availability authorizations. <input type="checkbox"/> is MODIFIED - Limitations for all documents processed in the STI system under the blanket release should be changed to conform to blocks as checked in Sections II and III.

I HAVE DETERMINED THAT THIS PUBLICATION:

- ☐ DOES contain export controlled, confidential commercial information, and/or discloses an invention for which a patent has been applied, and the appropriate limitation is checked in Sections III and/or IV.
- ☒ does NOT contain export controlled, confidential commercial information, nor does it disclose an invention for which a patent has been applied, and may be released as indicated above.

NAME OF AUTHOR/ORIGINATOR Rebecca Smith	MAIL CODE Dym-1	SIGNATURE <i>Rebecca S. Smith</i>	DATE 7 Apr 98
--------------------------------------------	--------------------	--------------------------------------	------------------

VII. PROJECT OFFICER/TECHNICAL MONITOR/DIVISION CHIEF REVIEW

- ☒ APPROVED FOR DISTRIBUTION AS MARKED ON REVERSE ☐ NOT APPROVED

NAME OF PROJECT OFFICER OR TECH. MONITOR RAYMOND M. WHEELER	MAIL CODE JT-G	SIGNATURE <i>Ray M. Wheeler</i>	DATE 4/10/98
----------------------------------------------------------------	-------------------	------------------------------------	-----------------

VIII. EXPORT CONTROL REVIEW/CONFIRMATION

- ☒ Public release is approved ☐ Export controlled limitation is not applicable
- ☐ Export controlled limitation is approved ☐ Export controlled limitation (ITAR/EAR) marked in Section III is assigned to this document:

USML CATEGORY NUMBER	CCL ECCN NUMBER	HQ OR CENTER EXPORT CONTROL ADMINISTRATOR (as applicable) <i>Dennis L. Miller</i>	DATE 4/16/98
----------------------	-----------------	--------------------------------------------------------------------------------------	-----------------

IX. PROGRAM OFFICE OR DELEGATED AUTHORITY REVIEW

- ☒ APPROVED FOR DISTRIBUTION AS MARKED ON REVERSE ☐ NOT APPROVED CODE M, DAA LETTER, 6/14/94

NAME OF PROGRAM OFFICE REPRESENTATIVE Audrey Lee Silipo	MAIL CODE FF-S2-A	SIGNATURE <i>Audrey Lee Silipo</i>	DATE 4/14/98
------------------------------------------------------------	----------------------	---------------------------------------	-----------------

X. DISPOSITION

THIS FORM, WHEN COMPLETED, IS TO BE SENT TO YOUR CENTER PUBLICATIONS OFFICE

INSTRUCTIONS FOR COMPLETING THE NASA SCIENTIFIC AND TECHNICAL DOCUMENT AVAILABILITY AUTHORIZATION (DAA) FORM

Purpose. This DAA form is used to prescribe the availability and distribution of all NASA-generated and NASA-funded documents containing scientific and technical information (including those distributed via electronic media such as the World Wide Web and CD-ROM).

Requirements. The author/originator must provide either a suitable summary description (title, abstract, etc.) or a completed copy of the document with this form. This form is initiated by the document author/originator and that individual is responsible for recommending/determining the availability/distribution of the document. The author/originator completes Sections I through III, and VI. The author/originator is also responsible for obtaining information and signature in Section IV to the extent the document discloses an invention for which patent protection has been applied. Subsequent to completion of these sections, the author/originator forwards the document to the appropriate Project Manager/Technical Monitor/Division Chief for further review and approval in Section VII, including a re-review of the planned availability and distribution. Once this approval is obtained, the DAA is forwarded to the NASA Headquarters or Center Export Administrator for completion of Section VIII. It is then forwarded for completion of Section IX to the cognizant NASA Headquarters Program Office or Delegated Authority, who provides final review and approval for release of the document as marked.

When to Use This Form. Documents containing STI and intended for presentation or publication (including via electronic media) must be approved in accordance with the NASA STI Procedures and Guidelines (NPG 2200.2). Documents that are to be published in the NASA STI Report Series must be coordinated with the appropriate NASA Headquarters or Center Scientific and Technical Information Office in accordance with NPG 2200.2. Note that information on the Report Documentation Page (if attached) is not to be entered on the DAA except for title, document date, and contract number.

How to Use this Form. Specific guidelines for each section of this form are detailed below.

I. Document/Project Identification. Provide the information requested. If the document is classified, provide instead the security classification of the title and abstract. (Classified information must not be entered on this form). Include RTOP numbers on the Contract/Grant/Interagency/Project Number(s) line. Provide information on presentations or externally published documents as applicable.

II. Security Classification. Enter the applicable security classification for the document. Documents, if classified, will be available only to appropriately cleared personnel having a "need to know."

III. Availability Category for Unclassified Documents. Check the appropriate category or categories.

Export Controlled Document. If the document is subject to export restrictions (see NPG 2200.2, paragraph 4.5.3), the appropriate restriction must be checked, either International Traffic in Arms Regulations (ITAR) or Export Administration Regulations (EAR), and the appropriate United States Munitions List (USML) category or Commerce Control List (CCL), Export Control Classification Number (ECCN) must be cited.

Confidential Commercial Documents (Documents containing Trade Secrets, SBIR documents, and/or Copyrighted Information). Check the applicable box (see NPG 2200.2 paragraph 4.5.7). When any of these boxes are checked, also indicate the appropriate limitation and expiration in the list to the right of these restrictions. These limitations refer to the user groups authorized to obtain the document. The limitations apply both to the initial distribution of the documents and the handling of requests for the documents. The limitations will appear on and apply to reproduced copies of the document. Documents limited to NASA personnel should not be made available to onsite contractors. If the Available Only With the Approval of Issuing Office limitation is checked, the NASA Center for Aerospace Information will provide only bibliographic processing and no initial distribution; CASI will refer all document requests to the Issuing office.

Publicly Available Document - Unrestricted Distribution. Check this box if the information in the document may be made available to the general public without restrictions (unrestricted domestic and international distribution). If the document is copyrighted (see paragraph 4.5.7.3 in NPG 2200.2), also check the "Copyrighted" box in this section.

IV. Document Disclosing an Invention. This must be completed when the document contains information that discloses an invention (see NPG 2200.2, paragraph 4.5.9). When this box is checked, an additional appropriate availability category must be checked. Use of this category must be approved by NASA Headquarters or Center Patent Counsel or the Intellectual Property Counsel.

V. Blanket Release (Optional). Complete this optional section whenever subsequent documents produced under the contract, grant, or project are to be given the same distribution and/or availability as described in Sections II and III. More than one contract number or RTOP Number can be entered. This section may also be used to rescind or modify an earlier Blanket Release. All blanket releases must be approved by the Program Office or its designee and concurred with by the Office of Management Systems and Facilities.

VI. Author/Originator Verification. Required for all DAA forms.

VII. Project Officer/Technical Monitor/Division Chief Review. The Project Officer/Technical Monitor/Author or Originator Division Chief or above must sign and date the form. The office code and typed name should be entered.

VIII. Export Control Review/Confirmation. This section is to be completed by the authorized NASA Headquarters or Center Export Control Administrator for all documents.

IX. Program Office or Delegated Authority Review. This section is to be completed by the duly authorized official representing the NASA Headquarters Program Office. Any delegation from NASA Headquarters to a NASA Center in accordance with NPG 2200.2 should be entered here.

X. Disposition. For NASA Center use.

Clinorotation Affects Morphology and Ethylene Production in Soybean Seedlings

Emmanuel Hilaire^{1,2}, Barbara V. Peterson¹, James A. Guikema² and Christopher S. Brown³

¹ Dynamac Corporation, Mail Code DYN-3, Kennedy Space Center, FL 32899, USA

² Kansas State University, Division of Biology, 232 Ackert Hall, Manhattan, KS 66506-4901, USA

³ Dynamac Corporation, 1910 Sedwick Rd., Building 100, Durham, NC 27713, USA

The microgravity environment of spaceflight influences growth, morphology and metabolism in etiolated germinating soybean. To determine if clinorotation will similarly impact these processes, we conducted ground-based studies in conjunction with two space experiment opportunities. Soybean (*Glycine max* [L.] Merr.) seeds were planted within BRIC (Biological Research In Canister) canisters and grown for seven days at 20°C under clinorotation (1 rpm) conditions or in a stationary upright mode. Gas samples were taken daily and plants were harvested after seven days for measurement of growth and morphology. Compared to the stationary upright controls, plants exposed to clinorotation exhibited increased root length (125% greater) and fresh weight (42% greater), whereas shoot length and fresh weight decreased by 33% and 16% respectively. Plants grown under clinorotation produced twice as much ethylene as the stationary controls. Seedlings treated with triiodo benzoic acid (TIBA), an auxin transport inhibitor, under clinorotation produced 50% less ethylene than the untreated control subjected to the same gravity treatment, whereas a treatment with 2,4-D increased ethylene by five-fold in the clinorotated plants. These data suggest that slow clinorotation influences biomass partitioning and ethylene production in etiolated soybean plants.

Key words: Auxin — Clinostat — Ethylene — *Glycine max* L. — Gravity — Partitioning.

Although the microgravity conditions of spaceflight do not appear to affect seed germination (for review see Dutcher et al. 1994, Claassen and Spooner 1994), it can influence subsequent plant growth. Brown et al. (1995) observed that soybean seedlings grown for 8 or 11 days in space had longer roots and higher root fresh weight than the ground controls. Cowles et al. (1984) found decreased growth (measured as plant length, fresh weight and dry weight) for pine, oat and mung bean plants developed in microgravity. Gallegos et al. (1995) noted that sweet clover seedlings were 40% longer relative to the ground controls

when grown for 3 days in space. Although these studies suggest that microgravity impacts growth by redirecting whole plant partitioning of assimilates, the mechanism is not clear.

The slowly rotating horizontal clinostat is a ground-based apparatus used to create a gravity vector-averaged environment and is sometimes useful to predict potential spaceflight effects on biological specimens (Albrecht-Buehler 1992). Several reports describe the effects of clinorotation on plant physiology such as cytoskeletal function (Hilaire et al. 1995a), protein expression (Piastruch and Brown 1995), carbohydrate metabolism (Brown and Piastruch 1994, Obenland and Brown 1994), calcium distribution (Hilaire et al. 1995b) and the cell cycle (Legué et al. 1992).

Ethylene, (CH₂)₂, is a flammable volatile plant hormone which diffuses in the gas phase through the intercellular spaces and outside the tissue (Jackson 1991). Under non-stress conditions, ethylene release occurs during different developmental stages of the plant such as fruit ripening and senescence (McAfee and Morgan 1971). There are many examples of increased ethylene production following stresses induced by abiotic or biotic agents. For example, plants exposed to supraoptimal temperatures (Field 1981) or flooding (Kawase 1976) produced increased amounts of ethylene. At the cellular level, ethylene has been shown to affect microtubule orientation (Lang et al. 1982), to increase cell wall thickness (Freytag et al. 1977), and to increase respiration and concentration of fructose 2,6-bisphosphate (Stitt et al. 1986). Several authors also reported an increase in ethylene production by plants exposed to clinorotation (Leather et al. 1972, Hensel and Iversen 1980, Driss-Ecole et al. 1994). Brown et al. (1995) extended these experiments to the microgravity environment and observed a two-fold increase in ethylene production in soybean seedlings grown in space relative to the ground controls.

The objectives of the present study were to determine (1) the influence of clinorotation on the growth, morphology and ethylene evolution of soybean seedlings and (2) if the addition of synthetic auxin or auxin transport inhibitors would affect the concentration of head space ethylene due to clinorotation. This study revealed an effect of clinorotation on soybean seedling morphology associated with an increase in ethylene production.

Abbreviations: BRIC, Biological Research In Canister; FAA, formaldehyde acetic acid; STS, Space Transportation System; TIBA, triiodo benzoic acid.




Fig. 1 (A) BRIC (Biological Research in Canister) ground support equipment hardware and (B) Teflon sleeve, germination paper seed germination apparatus.

Materials and Methods

Soybean (*Glycine max* L. [Merr] cv. McCall) seeds were germinated and grown under conditions of slow clinorotation (1 rpm) or maintained stationary for 7 days in BRIC (Biological Research in Canister) ground support hardware (Fig. 1A) at 20°C. These canisters consist of two independent compartments, each of which is 8.2 cm (diameter) by 31.2 cm (height). Each compartment contains 4 light-tight, passive pressure relief vents (1.5 mm in diameter) which were used as gas sampling ports. Individual seeds were rolled in germination paper, each roll was inserted into a Teflon tube (Fig. 1B) and each canister compartment contained 13 of such tubes. To initiate the experiment, 2.5 ml of sterile water, 10^{-4} M TIBA or 10^{-5} M 2,4-D was added to each roll. Duplicates gas samples (0.5 ml) were withdrawn daily from the head space of each canister compartment using gas-tight syringes. Analyses of the amounts of ethylene and CO_2 were performed immediately after sampling. Each gas sample was injected in a gas chromatograph equipped with a thermal detector for CO_2 (Hewlett Packard gas chromatograph 5880) or a photoionization detector for ethylene (Photovac gas chromatograph 10S70). The detector response was standardized against injections of known quantities of CO_2 or ethylene prepared by serial dilutions.

After the daily gas sampling, plants were removed from independent canisters and the shoots and roots were measured and weighed. The tissue was fixed in FAA and processed for light microscopy. Cross sectional areas of the shoot and root were analyzed by computer using the NIH Image 931220 software. Shoot sections were taken midway between the root/shoot interface and the hypocotyl hook. Root sections were taken midway between the root/shoot interface and the root tip.

Results

Soybean seedlings grown for 7 days on the horizontal clinostat at 1 rpm showed a decrease in shoot length by 33% (Fig. 2A) and a doubling of root length compared to the stationary controls (Fig. 2B). This difference in shoot and root length became evident by 4 to 5 days after planting. Shoots from the clinorotated plants were characterized by a decrease in fresh weight whereas roots from these plants had a greater fresh weight relative to the stationary controls (Table 1). Shoots from the clinorotated plants had a larger diameter relative to the stationary control whereas roots from the clinorotated plants had a smaller cross sectional area compared to the controls (Table 2).

Daily gas sampling from the canister revealed an increase in ethylene production through time in the stationary upright controls. By day 7 after planting, however, soybean seedlings grown under clinorotation conditions produced twice as much ethylene as the stationary control (Fig. 3A). The CO_2 concentration also increased through the plant development but no significant differences were found between the clinorotation treatment and the stationary control (Fig. 3B). When clinorotated seedlings were grown in the presence of an auxin transport blocker (TIBA), the amount of ethylene produced was two-fold lower starting at day 5 compared to the amount produced by clinorotated plants grown in water (Fig. 4A). The level of CO_2 produced by the TIBA treated plants was always higher than the control until day 7 where they were identical (Fig. 4B). On the other hand, treatment of the seedlings




Fig. 2 Soybean seedling growth in canisters under two conditions: upright stationary and horizontal clinorotation (1 rpm). Each symbol represents the mean value ($n=26-50 \pm \text{SE}$). Notice the decrease in shoot length (A) and the increase in root length resulting from clinorotation (B).

with 2,4-D during clinorotation resulted in a five-fold increase in ethylene production relative to the plants grown in water at day 5 after planting (Fig. 4A) and thereafter the level starts to decrease to return to the same level as the control in water only. The seedlings treated with 2,4-D produced more CO_2 relative to the control until day 7 where both levels were very close (Fig. 4B).

Table 1 Fresh weight of shoot and root tissue of seven-day-old etiolated soybean plants grown in stationary, upright position or exposed to clinorotation (1 rpm)

	Fresh weight (mg)		% Difference
	Stationary	Clinorotation	
Shoot	584 (24)	494 (24)*	-16%
Root	87 (8)	124 (8)*	+42%

Values represent the mean ($n=26$) and the standard error is shown in parentheses.

* Statistically different at the 5% level of significance.

Discussion

Clinorotation increases root growth relative to shoot growth in etiolated soybean seedlings, consistent with similar observations by Hoson et al. (1992) with azuki bean using a three-dimensional clinostat. These particular morphological changes have been observed previously in space-grown plants (Brown et al. 1995, Gallegos et al. 1995). Huber (1983) showed that starch stored in leaves during the day was degraded at night and preferentially translocated (as sucrose) to the root for growth. It has been shown that starch concentration in the cotyledons of clinorotated soybean seedlings is lower, perhaps indicative of diminished starch synthetic capability (Brown and Piastuch 1994). This suggests that the enhanced root growth under clinorotation may be related to the lower starch concentration due to enhanced carbon translocation from the cotyledons. Microscopical analysis showed that the effects of clinorotation on growth are associated with an increase in the diameter of the shoot and a decrease for the root.

Table 2 Cross sectional areas of stem and root tissue^a of seven-day-old etiolated soybean plants grown in stationary, upright position or exposed to clinorotation (1 rpm)

	Area (mm^2)		% Difference
	Stationary	Clinorotation	
Shoot	5.0 (0.3)	6.7 (0.5)*	+34%
Root	1.6 (0.2)	0.5 (0.1)*	-69%

Values represent the mean ($n=10$) and the standard error is shown in parentheses.

^a Shoot sections were taken midway between the root/shoot interface and the hypocotyl hook and root sections were taken midway between the root/shoot interface and the root tip.

* Statistically different at the 5% level of significance.







Fig. 3 (A) Ethylene and (B) CO₂ concentrations in the BRIC headspace with soybean seedlings grown under stationary conditions or under horizontal clinorotation (1 rpm). Each symbol represents the mean value ($n=15-16 \pm SE$).

Gas measurements revealed an increase in ethylene production during clinorotation. Taken together, these data suggest that the increase in ethylene due to clinorotation may affect root and shoot tissue morphology. Leather et al. (1972) reported that 8-week-old tomato plants produced twice as much ethylene after two hours of clinorotation at 0.2 rpm relative to the controls. This rise in ethylene con-

Fig. 4 (A) Ethylene (B) CO₂ concentrations in the BRIC headspace with soybean seedlings grown under horizontal clinorotation (1 rpm) in the presence of water, 10^{-4} M TIBA or 10^{-5} M 2,4D. Each symbol represents the mean value ($n=15-16 \pm SE$).

centration was associated with leaf epinasty. Hensel and Iversen (1980) observed that 24-h-old cress seedlings produced at least three times more ethylene than the stationary controls after five hours of clinorotation at 2 rpm. The same authors also found that the root length of the clinorotated seedlings was 81.4% of that of the controls.

Both of these experiments described a rise in ethylene within hours after clinorotation whereas we observed a similar increase after seven days of growth and clinorotation, indicating that the clinorotation-induced ethylene production is not a short-term transient effect. This is corroborated by a more recent study from Driss-Ecole et al. (1994) who grew *Veronica arvensis* plants for 45 days on a clinostat rotating at 1 rpm and found that ethylene production was higher in the clinorotated plants relative to the rotating vertical controls. However, when ethylene concentration was expressed relative to plant biomass no differences were found. These authors observed that the hypocotyl and root dry weight from the clinorotated samples was increased by 40% relative to the controls, explained by an increase in secondary root production. In our experiment, 7-day-old clinorotated soybean seedlings had longer main roots than the stationary controls but very little secondary roots. If the experiment would have continued for a longer period of time, it is possible that enhanced secondary root growth would have occurred.

Root elongation has been shown to be promoted (Dedolph et al. 1965, Smith and Robertson 1971, Koning and Jackson 1979, Nakayama and Ota 1980) or inhibited by ethylene (Chadwick and Burg 1967, Rauser and Horton 1975, Whalen and Feldman 1988), while other reports suggest an ethylene-induced decrease in shoot growth (Warner and Leopold 1971, Goeschl and Kays 1975, Nee et al. 1978). Therefore, it is possible that the alteration in root/shoot ratios which we observed upon clinorotation are due in part to an ethylene buildup. We observed that ethylene production was significantly higher than the stationary control at day 3 after planting. Since at that age the seedlings are still very short (about 30 mm) and therefore are still firmly maintained within the germination paper, it is likely that this increase in ethylene production is due to a disturbance in gravity perception produced by clinorotation rather than to mechanical stress, such as the contact of the seedlings with the teflon sleeve. In agreement with our results is the report from Salisbury and Wheeler (1981) who showed that leaf epinasty observed in clinorotated plants was due to ethylene production and that epinasty was caused by disturbances in the gravity perception mechanism rather than leaf flopping.

Previous studies reported that high ethylene concentration causes seedling swelling (Bertell et al. 1990, Camp and Wicklife 1981) by affecting microtubule and cellulose microfibril orientation (Eisinger 1983), therefore altering seedling growth (Smith and Robertson 1971). Clinorotation did not alter CO₂ concentration significantly compared to the stationary controls. Previous studies from spaceflight experiments also report an increase in ethylene production in microgravity but no changes in CO₂ (Brown et al. 1995, Gallegos et al. 1995) suggesting that both horizontal slow clinorotation and microgravity represent stressful en-

vironments for the plants. However, in both of these space experiments gas samples were taken after the seedlings were back to Earth, therefore the increase in ethylene production may be due to the stress produced by shuttle re-entry in the atmosphere and/or to re-exposure to Earth unit gravity and not to microgravity. Caution must be taken in interpreting the results because even if an increase in ethylene production was observed in clinorotated samples and in some space flight samples (Brown et al. 1995, Gallegos et al. 1995), it does not necessarily imply that the cellular mechanism leading to this result is identical. In the case of clinorotation, the constant movement of the amyloplasts within the columella cells (Hilaire et al. 1995a) may constitute a cellular mechanical stress (as opposed of a physical contact which would be an external mechanical stress) whereas in the case of space environment it would be the reduced gravity. In both conditions, the stress is constant as well as the increase in ethylene production. In addition, it is important to note that ethylene has been proposed to be required in stem gravitropism (Wheeler and Salisbury 1980). Therefore, during gravistimulation the ethylene production may be enhanced in a transient manner and correlated with the generation of a curvature, possibly due to the rapid movement of the amyloplasts from one region of the columella cells to the new distal pole.

When treated with an auxin transport inhibitor (TIBA) or a synthetic auxin analogue (2,4-D) during clinorotation, soybean seedlings produced more CO₂ than the control and reached a plateau faster. In addition, the TIBA treatment during clinorotation reduced the production of ethylene by the plants whereas addition of 2,4-D increased ethylene concentration considerably. Exogenously added 2,4-D is subsequently degraded (Kang et al. 1971) and presumably is no longer effective, which could explain the drop in ethylene concentration observed after day 5.

Previous studies have explored the relationship between auxin and ethylene production. For example, Romano et al. (1993) reported that overproduction of auxin in transgenic plants resulted in the overproduction of ethylene. In addition, Yoshii and Imaseki (1981) showed that auxin promotes ethylene production by inducing the synthesis of ACC synthase, an enzyme involved in the biosynthesis and regulation of ethylene (Yang and Hoffman 1984). Furthermore, Dedolph et al. (1966) observed that the auxin sensitivity of clinorotated *Avena* seedlings to applied IAA was increased and a recent study from Li and Wu (1995) reported that the expression of a GUS reporter fused to an auxin-induced promoter sequence (SAUR) was enhanced in *Arabidopsis* seedlings transformed with this construct grown in microgravity or under clinorotation conditions. We believe that clinorotation creates an auxin imbalance within the soybean seedlings and therefore a high concentration of auxin within a specific region of the seedlings would induce ACC synthase and in turn ethylene pro-

duction. This increase in ethylene would decrease shoot growth and increase root elongation. In the limit of our study, we cannot predict what tissue has the highest concentration of auxin nor if cell sensitivity to auxin is affected by clinorotation.

The authors wish to thank Dr. Yi Li for his valuable comments. Work was conducted under NASA contract NAS10-12180 to Dynamac Corporation. This represents contribution 96-556-J of the Kansas Agricultural Experiment Station.

References

- Albrecht-Buehler, A. (1992) The simulation of microgravity conditions on the ground. *ASGS Bull.* 5: 3-10.
- Bertell, G., Bolander, E. and Eliasson, L. (1990) Factors increasing ethylene production enhance the sensitivity of root growth to auxins. *Physiol. Plant.* 79: 255-258.
- Brown, C.S., Hilaire, E.M., Guikema, J.A., Piastuch, W.C., Johnson, C.F., Stryjewski, E.C., Peterson, B.V. and Vordermark, D.S. (1995) Metabolism, ultrastructure and growth of soybean seedlings in microgravity: results from the BRIC-01 and BRIC-03 experiments. *ASGS Bull.* 9: 93.
- Brown, C.S. and Piastuch, W.C. (1994) Starch metabolism in germinating soybean cotyledons is sensitive to clinorotation. *Plant Cell Environ.* 17: 341-344.
- Camp, P.J. and Wicklife, J.F. (1981) Light or ethylene treatments induce cell enlargement in etiolated maize mesocotyls. *Plant Physiol.* 67: 125-128.
- Chadwick, A.V. and Burg, S.P. (1967) An explanation of the inhibition of root growth caused by indole-3-acetic acid. *Plant Physiol.* 42: 415-420.
- Claassen, D.E. and Spooner, B.S. (1994) Impact of altered gravity on aspects of cell biology. *Int. Rev. Cytol.* 156: 301-373.
- Cowles, J.R., Scheld, H.W., Lemay, R. and Peterson, C. (1984) Growth and lignification in seedlings exposed to eight days of microgravity. *Ann. Bot.* 54: 33-48.
- Dedolph, R.R., Naqvi, S.M. and Gordon, S.A. (1965) Effect of gravity compensation on the geotropic sensitivity of *Avena* seedlings. *Plant Physiol.* 40: 961-965.
- Dedolph, R.R., Naqvi, S.M. and Gordon, S.A. (1966) Role of indole-3-acetic acid in modification of geotropic responses in clinostat rotated *Avena* seedlings. *Plant Physiol.* 41: 897-902.
- Driss-Ecole, D., Cottignies, A., Jeune, B., Corbineau, F. and Perbal, G. (1994) Increased mass production of *Veronica arvensis* grown on a slowly rotating clinostat. *Environ. Exp. Bot.* 34: 303-310.
- Dutcher, F.R., Hess, E.L. and Halstead, T.W. (1994) Progress in plant research in space. *Adv. Space Res.* 14: 159-171.
- Eisinger, W. (1983) Regulation of pea internode expansion by ethylene. *Annu. Rev. Plant Physiol.* 34: 225-240.
- Field, R.J. (1981) A relationship between membrane permeability and ethylene production at high temperature in leaf tissue of *Phaseolus vulgaris* L. *Ann. Bot.* 48: 33-39.
- Freytag, A.H., Berlin, J.D. and Linden, J.C. (1977) Ethylene-induced fine structure alterations in cotton and sugarbeet radicle cells. *Plant Physiol.* 60: 140-143.
- Gallegos, G.L., Odum, W.R. and Guikema, J.A. (1995) Effect of microgravity on stress ethylene and carbon dioxide production in sweet clover (*Mellilotus alba* L.). *J. Grav. Physiol.* 2: 155-156.
- Goeschl, J.D. and Kays, S.J. (1975) Concentration dependencies of some effects of ethylene on etiolated pea, peanut, bean and cotton seedlings. *Plant Physiol.* 55: 670-677.
- Hensel, W. and Iversen, T.H. (1980) Ethylene production during clinostat rotation and effect on root geotropism. *Z. Pflanzenphysiol.* 97: 343-348.
- Hilaire, E., Paulsen, A.Q., Brown, C.S. and Guikema, J.A. (1995a) Effects of clinorotation and microgravity on sweet clover columella cells treated with cytochalasin D. *Physiol. Plant.* 95: 267-273.
- Hilaire, E., Paulsen, A.Q., Brown, C.S. and Guikema, J.A. (1995b) Microgravity and clinorotation cause redistribution of free calcium in sweet clover columella cells. *Plant Cell Physiol.* 36: 831-837.
- Hosono, T., Kamisaka, S., Masuda, Y. and Yamashita, M. (1992) Changes in plant growth processes under microgravity conditions simulated by a three-dimensional clinostat. *Bot. Mag.* 105: 53-70.
- Huber, S.C. (1983) Relation between photosynthetic starch formation and dry weight partitioning between the root and shoot. *Can. J. Bot.* 61: 2709-2716.
- Jackson, M.B. (1991) Ethylene in root growth and development. In *The Plant Hormone Ethylene*. Edited by Mattoo, A.K. and Suttle, J.C. pp. 159-182. CRC Press, Boca Raton, FL.
- Kang, B.G., Newcomb, W. and Burg, S.P. (1971) Mechanism of auxin-induced ethylene production. *Plant Physiol.* 47: 504-509.
- Kawase, M. (1976) Ethylene accumulation in flooded plants. *Physiol. Plant.* 36: 236-241.
- Konings, H. and Jackson, M.B. (1979) A relationship between the rates of ethylene production by roots and the promoting or inhibiting effects of exogenous ethylene and water on root elongation. *Z. Pflanzenphysiol.* 92: 385-397.
- Lang, J.M., Eisinger, W.R. and Green, P.B. (1982) Effects of ethylene on the orientation of microtubules and cellulose microfibrils of pea epicotyls with polyamellate cell walls. *Protoplasma* 110: 5-14.
- Leather, G.R., Forrence, L.E. and Abeles, F.B. (1972) Increase ethylene production during clinostat experiments may cause leaf epinasty. *Plant Physiol.* 49: 183-186.
- Legué, V., Driss-Ecole, D. and Perbal, G. (1992) Cell cycle and cell differentiation in lentil roots grown on a slowly rotating clinostat. *Physiol. Plant.* 84: 386-392.
- Li, Y. and Wu, Y. (1995) Effects of clinorotation and microgravity on expression of auxin-inducible SAUR and GH3 genes in transgenic plants. *ASGS Bull.* 9: 54.
- McAfee, J.A. and Morgan, P.W. (1971) Rates of production and internal levels of ethylene in vegetative cotton plant. *Plant Cell Physiol.* 12: 839-847.
- Nakayama, M. and Ota, Y. (1980) Physiological action of ethylene in crop plants. VI. Effect of hydrocarbons especially ethylene on the root growth of soybean and rice seedlings. *Japan J. Crop Sci.* 49: 360-372.
- Nee, M., Chiu, L. and Eisinger, W. (1978) Induction of swelling in pea internode tissue by ethylene. *Plant Physiol.* 62: 902-906.
- Obenland, D.M. and Brown, C.S. (1994) The influence of altered gravity on carbohydrate metabolism in excised wheat leaves. *J. Plant Physiol.* 144: 696-699.
- Piastuch, W.C. and Brown, C.S. (1995) Protein expression in *Arabidopsis thaliana* after chronic clinorotation. *J. Plant Physiol.* 146: 329-332.
- Rausser, W.E. and Horton, R.F. (1975) Rapid effects of indoleacetic acid and ethylene on the growth of intact pea roots. *Plant Physiol.* 55: 443-447.
- Romano, C.P., Cooper, M.L. and Klee, H.J. (1993) Uncoupling auxin and ethylene effects in transgenic tobacco and arabidopsis plants. *Plant Cell* 5: 181-189.
- Salisbury, F.B. and Wheeler, R.M. (1981) Interpreting plant responses to clinostating: I. Mechanical stresses and ethylene. *Plant Physiol.* 67: 677-685.
- Smith, K.A. and Robertson, P.D. (1971) Effect of ethylene on root extension of cereals. *Nature* 234: 148-149.
- Stitt, M., Cseke, C. and Buchanan, B. (1986) Ethylene-induced increase in Fructose-2,6-bisphosphate in storage tissues. *Plant Physiol.* 80: 246-248.
- Warner, H.L. and Leopold, A.C. (1971) Timing of growth regulator responses in pea. *Biochem. Biophys. Res. Commun.* 44: 989-994.
- Whalen, M.C. and Feldman, L.J. (1988) The effect of ethylene on root growth of *Zea Mays* seedlings. *Can. J. Bot.* 66: 719-723.
- Wheeler, R.M. and Salisbury, F.B. (1980) Gravitropism in plant stems may require ethylene. *Science* 209: 1126-1128.
- Yang, S.F. and Hoffman, N.E. (1984) Ethylene biosynthesis and its regulation in higher plants. *Annu. Rev. Plant Physiol.* 35: 155-189.
- Yoshii, H. and Imaseki, H. (1981) Biosynthesis of auxin-induced ethylene. Effects of indole-3-acetic acid, benzyladenine and abscisic acid on endogenous levels of 1-aminocyclo-propane-1-carboxylic acid (ACC) and ACC synthase. *Plant Cell Physiol.* 22: 369-379.

(Received December 16, 1995; Accepted July 11, 1996)